


Spearhead*



Contains 20 g/l (1.72% w/w) clopyralid, 15.9 g/l (1.45% w/w) diflufenican and 300 g/l (25.9% w/w) MCPA.

- ★ **Season long control**
- ★ **Controls problem weeds like speedwell and clovers**
- ★ **Can be applied to young turf**
- ★ **Broad spectrum, residual activity**



White Clover



Slender Speedwell



Daisy

Weeds Controlled

SPEARHEAD is a selective herbicide for the control of a wide variety of broad leaved weeds in established turf. These include Speedwell, Daisy, Clover, Buttercup and Dandelion (see Application rates for additional weeds controlled).

Applications made under or followed by adverse conditions such as waterlogged and/or cold conditions may result in transient damage. Do not overlap spray swaths.

Timing of Application

For best results, the grass and weeds should be actively growing at the time of application. In most years, treatment during the early part of the season is recommended although spraying at any time between April and October will be effective, except during drought or waterlogged conditions. Only use SPEARHEAD on turf when the sward is satisfactorily established and regular mowing has begun under normal good management. **Young turf in the spring or early summer may be ready for spraying approximately two months after germination,** but turf sown in the late summer or autumn should not be sprayed until growth is resumed in the following spring, perhaps eight months after sowing.

Mowing

Avoid mowing during the 3 - 4 days preceding or following treatment. **CUTTINGS FROM THE TREATED AREA MUST NOT BE USED AS A MULCH FOR SOFT FRUIT, VEGETABLES, ORNAMENTALS OR ANY OTHER CROP.**

Drift

Avoid drift as much as possible by keeping the nozzles at the correct height and working only under conditions of little or no wind. Small amounts of SPEARHEAD spray may cause serious injury to herbaceous plants and fruit and vegetable crops.

Equipment

VEHICLE MOUNTED HYDRAULIC SPRAYER

VOLUME		
AREA	SPEARHEAD	WATER
one hectare	4 - 5 litres	200 - 700 litres

KNAPSACK SPRAYER

VOLUME		
AREA	SPEARHEAD	WATER
100 m ²	40 - 50 ml	4.5 litres

Application rates

Use SPEARHEAD at 4 - 5 litres per hectare. Half fill sprayer with clean water. Add the required quantity of SPEARHEAD and complete filling. This aids the effective mixing for all Selective Herbicides.

4 litres per hectare

Creeping Buttercup, Greater Plantain, Ribwort Plantain, White Clover

5 litres per hectare

Autumn Hawkbit
 Black Medick (from seed - up to 2 expanded true leaves)
 Charlock (up to 6 expanded true leaves)
 Common chickweed*
 Common Mouse-ear
 Corn Marigold (up to 6 expanded true leaves)

PROTECT FROM FROST.

ALWAYS READ THE LABEL. USE PESTICIDES SAFELY.
 (MAPP 09941)

PACK COVERAGE: 5 LITRES COVERS 10,000 m²

*Spearhead is a Trademark of Dow AgroSciences.
 Clopyralid is supplied by Dow AgroSciences.

Full product information, label text and MSDS are available for download from www.bayer-escience.co.uk

Daisy
 Dandelion
 Mayweeds (up to 6 expanded true leaves)
 Mouse-ear Hawkweed
 Self Heal*
 Slender Speedwell*
 Smooth Sowthistle (up to 6 expanded true leaves)
 Sorrels*
 Thrift
 Trefoils (up to 2 expanded true leaves)
 Vetches (from seed - up to 2 expanded true leaves)
 Yarrow*
 Yellow Suckling Clover

*Moderate control. Complete control may not always be achieved.

Qualified Minor Use Recommendation
 Limited evidence suggests that the following weeds may be susceptible to SPEARHEAD at the 5L/ha dose:

Black bindweed (up to 2 expanded true leaves)
 Creeping Thistle (from seed - up to 6 expanded true leaves)
 Groundsel (up to 6 expanded true leaves)

After use

Wash equipment thoroughly after use.

